

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 956 854 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

17.11.1999 Bulletin 1999/46

(51) Int. Cl.⁶: **A61K 31/12**, **A61K 31/05**(21) Application number: **97935798.5**

(86) International application number:

PCT/JP97/02845(22) Date of filing: **18.08.1997**

(87) International publication number:

WO 98/07417 (26.02.1998 Gazette 1998/08)

(84) Designated Contracting States:

BE CH DE DK ES FR GB IT LI NL SE(30) Priority: **16.08.1996 JP 23472996****13.06.1997 JP 17319197**(71) Applicant: **KANEKA CORPORATION****Osaka-shi, Osaka 530-8288 (JP)**

(72) Inventors:

• **MAE, Tatsumasa****Kakogawa-shi, Hyogo 675-01 (JP)**• **SAKAMOTO, Yoshitomo****Akashi-shi, Hyogo 674 (JP)**• **MORIKAWA, Souichi****Himeji-shi, Hyogo 670 (JP)**• **HIDAKA, Takayoshi****Kobe-shi, Hyogo 655 (JP)**

(74) Representative:

Hubert, Philippe et al**Cabinet Beau de Loménie****158, rue de l'Université****75340 Paris Cédex 07 (FR)****(54) PHARMACEUTICAL COMPOSITION COMPRISING COENZYME Q 10**

(57) The present invention has for its object to provide a medicinal composition comprising coenzyme Q₁₀ as an active ingredient, which composition features an enhanced absorption after oral administration.

The present invention is directed to a medicinal composition comprising coenzyme Q₁₀ as an active ingredient with the reduced form of coenzyme Q₁₀ accounting for more than 20 weight % of said coenzyme Q₁₀.


EP 0 956 854 A1

EP 0 956 854 A1

Description

TECHNICAL FIELD

- 5 [0001] The present invention relates to a medicinal composition with improved absorption after oral administration which comprises a coenzyme Q₁₀ of the following general formula (1-A) as an active ingredient.



BACKGROUND ART

- 25 [0002] Coenzyme Q₁₀ is a class of physiological substances occurring as component factors of the mitochondrial electron transfer system within the biological cell. Coenzyme Q₁₀ acts directly as an electron carrier in the oxidative phosphorylation reactions, through metabolic pathways, particularly aerobic pathways, to produce ATP and hence energy.
- 30 [0003] It seems that the demand for coenzyme Q₁₀ is increased in normal subjects in the state of severe physical fatigue and patients with cardiovascular disease, chronic debilitating disease, or on prolonged pharmacotherapy. It has been shown that a deficiency of coenzyme Q₁₀ occurs particularly in ischemic heart diseases, senile myocardial sclerosis, and hypertensive heart diseases. Therefore, it is a sound therapeutic choice to administer coenzyme Q₁₀ to those patients.
- 35 [0004] Moreover, coenzyme Q₁₀ has been used for non-therapeutic purposes as a nutrient or nutritional supplement just like vitamins.
- [0005] In order that coenzyme Q₁₀ may express its therapeutic efficacy or nutritional effect, it is most important to increase the coenzyme Q₁₀ level within the patient's tissue cells.
- 40 [0006] Coenzyme Q₁₀ is a lipid-soluble and practically water-insoluble substance and, therefore, it is only sparingly soluble in gastric juice. Consequently, oral dosage forms containing coenzyme Q₁₀ in solid state, such as tablets, granules, capsules, and suspensions for extemporaneous preparation, are not well absorbed after oral administration. This means that a considerably greater amount of coenzyme Q₁₀ than actually needed must be administered to the patient but such a practice tends to cause adverse gastrointestinal reactions such as epigastric discomfort, anorexia, nausea, and diarrheas.
- 45 [0007] Much research has heretofore been undertaken for overcoming those disadvantages. Japanese Kokai Publications Sho-55-81813 and Sho-61-221131, among others, disclose coenzyme Q₁₀ formulations of the solution type or the emulsion/dispersion type. However, such pharmaceutical devices are not sufficient to improve the absorption of coenzyme Q₁₀ in a satisfactory measure.
- 50 [0008] Japanese Kokai Publication Sho-56-18914 discloses a technology for accelerating the lymphatic absorption of coenzyme Q₁₀. This technology has been demonstrated to increase the absorption of coenzyme Q₁₀ in a certain measure but has not proved practically useful as yet.
- [0009] Japanese Kokai Publication Sho-60-89442 discloses a cyclodextrin-clathrated coenzyme Q₁₀ formulation. Japanese Kokai Publication Sho-60-1124 discloses a coenzyme Q₁₀-containing ribosomal formulation. However, those coenzyme Q₁₀ preparations require a complicated pharmaceutical procedure for production and are not practically fully
- 55 satisfactory.
- [0010] Italian Patent 1190442 Specification discloses a technology which, instead of using coenzyme Q₁₀ as such, comprises converting a reduced form of coenzyme Q₁₀ to a derivative such as an acyl ester, a sulfuric acid ester, or a phosphoric acid ester and administering this coenzyme Q₁₀ derivative for enhanced absorption. However, the effect of

EP 0 956 854 A1

the technology has not been supported by experimental data.

SUMMARY OF THE INVENTION

[0011] The present invention has for its object to provide a medicinal composition comprising coenzyme Q₁₀ as an active ingredient, which composition features an enhanced absorption after oral administration.

[0012] In the course of their intensive research for overcoming the above-mentioned disadvantages of the prior art, the inventors of the present invention discovered that when a medicinal composition containing a reduced form of coenzyme Q₁₀ was prepared and administered to patients by the oral route, a considerably higher bioavailability was surprisingly obtained as compared with the conventional medicinal composition containing only the oxidized form of coenzyme Q₁₀. The present invention has been developed on the basis of the above finding.

[0013] The present invention, therefore, is directed to a medicinal composition comprising coenzyme Q₁₀ as an active ingredient with the reduced form of coenzyme Q₁₀ accounting for more than 20 weight % of said coenzyme Q₁₀.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014]

Fig. 1 is a graphical representation of the relationship of plasma total coenzyme Q₁₀ concentration with the time after administration. The ordinate represents the plasma total coenzyme Q₁₀ concentration and the abscissa represents the time after administration. Each plot represents mean \pm standard deviation (n=4).


Fig. 2 is a graphical representation of the relationship of the total plasma coenzyme Q₁₀ concentration at 3 hours after administration with the weight ratio of oxidized form of coenzyme Q₁₀ to reduced form of coenzyme Q₁₀ in each sample. The ordinate represents the plasma total coenzyme Q₁₀ concentration and the abscissa represents the oxidized form of coenzyme Q₁₀-reduced form of coenzyme Q₁₀ ratio by weight. Each bar represents mean \pm standard deviation (n=4).

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention is now described in detail.

[0016] It is known that a fairly high proportion, usually about 40 to 90%, of coenzyme Q₁₀ occurs in reduced form in the body.

[0017] In vivo, the reduced form of coenzyme Q₁₀ is readily transformed to the oxidized form, while the oxidized form of coenzyme Q₁₀ is readily transformed into the reduced form. Therefore, coenzyme Q₁₀ in vivo can be generally expressed by the following formula (1).



[0018] Referring to the above formula (1), the general formula (1-A) represents the oxidized form of coenzyme Q₁₀ and the general formula (1-B) represents the reduced form of coenzyme Q₁₀.

[0019] In the conventional medicinal composition containing a coenzyme Q₁₀ as an active ingredient, the sole active ingredient is the oxidized form of coenzyme Q₁₀ of the above chemical formula (1-A). In contrast, the medicinal composition of the present invention comprises a reduced form of coenzyme Q₁₀ of the above chemical formula (1-B) as an active ingredient coenzyme Q₁₀. Consequently, as compared with the conventional medicinal composition containing only the oxidized form of coenzyme Q₁₀ as an active ingredient, the medicinal composition of the present invention is improved in absorption after oral administration and insures a higher bioavailability.

[0020] There is no particular limitation on the technology for providing said reduced form of coenzyme Q₁₀. A typical

EP 0 936 854 A1

method, which is by no means exclusive, comprises harvesting a coenzyme Q_{10} from a synthetic reaction mixture, a fermentation broth, or a natural source by procedures known in the art and subjecting it to chromatography to separate and concentrate the reduced form of coenzyme Q_{10} fraction. Where necessary, there can be followed the procedure of adding a conventional reducing agent such as sodium borohydride or sodium dithionite (sodium hydrosulfite) to the above coenzyme Q_{10} to reduce the oxidized form of coenzyme Q_{10} fraction of said coenzyme Q_{10} and, then, concentrate the reduced Q_{10} by chromatography. As a further alternative, the objective reduced form of coenzyme Q_{10} can be obtained by permitting said reducing agent to act on the available high-purity coenzyme Q_{10} .

[0021] There is no particular limitation on the technology for manufacturing the medicinal composition of the present invention. A typical but by no means exclusive method comprises dissolving the reduced form of coenzyme Q_{10} thus obtained and a commercial oxidized form of coenzyme Q_{10} in a suitable common solvent such as isopropyl alcohol, acetone, or ether to provide a medicinal composition containing said reduced form of coenzyme Q_{10} in a desired proportion. As an alternative, the above-mentioned reduced and oxidized forms of coenzyme Q_{10} can be simply admixed in solid stage. It is also possible to directly use the mixture of oxidized and reduced forms of coenzyme Q_{10} obtained in the course of the above-mentioned production process for coenzyme Q_{10} . Furthermore, the active ingredient for the medicinal composition of the present invention can be directly obtained by controlling the time of reduction reaction of the high-purity coenzyme Q_{10} already available and the type or amount of reducing agent to be used.

[0022] In the medicinal composition of the present invention, the reduced form of coenzyme Q_{10} accounts for more than 20 weight % of the total amount of coenzyme Q_{10} . If its proportion is not less than 20 weight %, the bioavailability of the resulting medicinal composition will not be as high as expected. The preferred proportion is not less than 40 weight % and the most preferred proportion is not less than 60 weight %. Conversely if the proportion of the reduced form of coenzyme Q_{10} is too large, the production process will be complicated and the cost of production increased. Therefore, it is not necessary to increase the coenzyme Q_{10} content too much.

[0023] The medicinal composition of the present invention can be used as, for example, a cardiogenic effective against symptoms in ischemic heart disease, senile myocardial sclerosis, hypertensive heart disease, etc. It can also be used as a nutrient, a nutritional supplement, or a veterinary medicine.

[0024] There is no particular limitation on the dosage form for the medicinal composition of the present invention. It may for example be powders, granules containing a binder component, or compression-molded tablets. Such powders or granules may be filled in capsule shells to provide capsules. They may also be processed into soft capsules by adding a natural oil, an oily higher fatty acid, a higher fatty acid monoglyceride, or a mixture thereof and wrapping the medicated oil in soft capsule sheet materials. In this application, the capsule shell may be one predominantly composed of gelatin or any other water-soluble macromolecular substance. The capsule includes microcapsules.

[0025] The medicinal composition of the present invention may contain, in addition to said reduced form of coenzyme Q_{10} , a variety of pharmaceutically acceptable formulating substances as added in suitable amounts in the routine manner. There is no particular limitation on the kinds of such substances. Thus, an excipient, a disintegrator, a lubricant, a binder, an antioxidant, a coloring agent, an antiflocculant, an absorption promoter, a solubilizer for the active ingredient, a stabilizer, etc. can be added as necessary.

[0026] The above-mentioned excipient includes but is not limited to sucrose, lactose, glucose, corn starch, mannitol, crystalline cellulose, calcium phosphate, and calcium sulfate.

[0027] The disintegrator includes but is not limited to starch, agar, calcium citrate, calcium carbonate, sodium hydrogen carbonate, dextrin, crystalline cellulose, carboxymethylcellulose, and gum tragacanth.

[0028] The lubricant includes but is not limited to talc, magnesium stearate, polyethylene glycol, silica, and hydrogenated vegetable oil.

[0029] The binder includes but is not limited to ethylcellulose, methylcellulose, hydroxypropylmethylcellulose, gum tragacanth, shellac, gelatin, gum arabic, polyvinylpyrrolidone, polyvinyl alcohol, polyacrylic acid, polymethacrylic acid, and sorbitol.

[0030] The antioxidant includes but is not limited to ascorbic acid, tocopherol, vitamin A, β -carotene, sodium hydrosulfite, sodium thiosulfate, sodium pyrosulfite, and citric acid.

[0031] There is no particular limitation on the coloring agent that can be used. For example, a variety of pharmaceutically acceptable colors can be mentioned.

[0032] The antiflocculant includes but is not limited to stearic acid, talc, light silicic anhydride, and hydrous silicon dioxide.

[0033] The absorption promoter includes but is not limited to higher alcohols, higher fatty acids, and glycerin fatty acid esters and other surfactants.

[0034] The above-mentioned solubilizer for the active ingredient includes but is not limited to organic acids such as fumaric acid, succinic acid, and malic acid.

[0035] The stabilizer includes but is not limited to benzoic acid, sodium benzoate, and ethyl p-hydroxybenzoate.

[0036] When the preparation comprising the medicinal composition of the present invention is used for by oral administration, the dosage should be selected according to the usage such as a drug, a veterinary medicine, or a nutrient.

EP 0 956 854 A1

[0037] For oral administration to domestic animals or fowls, the composition can be used as admixed into the feed or administered by a conventional forced manner.

BEST MODE FOR CARRYING OUT THE INVENTION

[0038] The following examples and formulation examples are intended to illustrate the present invention in further detail and should by no means be limitative of the scope of the present invention.

Example 1

(1) Preparation of samples

Preparation of Sample 1

[0039] A 5:95 (w/w) mixture (0.3 g) of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀ was melted on a water bath at 50°C and an olive oil was added to the melt to make 6.0 ml. This mixture was homogenized at 50°C to provide an oily composition.

Preparation of Comparative Sample 1

[0040] Oxidized coenzyme Q₁₀ (0.3 g) was melted on a water bath at 50°C and an olive oil was added to the melt to make 6.0 ml. This mixture was homogenized at 50°C to provide an oily composition.

(2) Oral absorption test

[0041] Sample 1 and Comparative Sample 1 were used as test samples. The test was performed using male Crj:CD (SD) rats (body weights 260 to 300 g) under well-fed conditions. As to dosage, each test sample was administered orally at the rate of 100 mg of total coenzyme Q₁₀ per kg body weight. In the test, the total plasma coenzyme Q₁₀ concentration was determined before administration (not administered) and serially after administration. Four rats were used per test sample for each time-point. The total coenzyme Q₁₀ means the sum of the mixture comprising the oxidized and reduced forms of coenzyme Q₁₀. The total plasma coenzyme Q₁₀ concentration was assayed as the concentration of oxidized form of coenzyme Q₁₀ in the following manner. To 1.0 ml of the obtained plasma sample, 2.0 ml of water, 4.0 ml of ethanol, and 10.0 ml of n-hexane were added in the order mentioned. The mixture was shaken vigorously for about 5 minutes and then centrifuged to separate into two layers. The organic layer was taken and the aqueous layer was further extracted with 10.0 ml of n-hexane twice in the same manner. The resulting organic layers and the organic layer previously taken were combined and evaporated to dryness. To the residue was added 250 µl of ethanol:1N-hydrochloric acid (99:1, v/v) for use as an assay sample. The assay of coenzyme Q₁₀ was carried out by high-performance liquid chromatography under the following conditions.

Column: 250 mm long x 4.6 mm in diameter SYMMETRY C18 (Waters)
Mobile phase: 0.5 M NaClO₄/C₂H₅OH:CH₃OH:CH₃CN:70%HClO₄ (400:300:300:1, v:v)
Detection wavelength: 275 nm
Flow rate: 1 ml/min.

[0042] The test results are presented in Fig. 1. In Fig. 1, the ordinate represents total plasma coenzyme Q₁₀ concentration and the abscissa represents the time after administration. Each plot is mean ± standard deviation.

[0043] It is apparent from Fig. 1 that whereas the plasma concentration peak appeared at 3 hr after administration in the case of the composition containing only the oxidized form of coenzyme Q₁₀, the peak appeared 1 hour earlier, i.e. at 2 hours after administration, in the case of the composition comprising the reduced form of coenzyme Q₁₀. Furthermore, the concentration level is also 2.1 times as high for the composition comprising the reduced form of coenzyme Q₁₀. It is, thus, clear that compared with the composition containing only the oxidized form of coenzyme Q₁₀, the medicinal composition of the present invention is absorbed faster and in a larger amount.

EP 0 956 854 A1

Example 2

(1) Preparation of samples

5 Preparation of Sample 2

[0044] Using a 20:80 (w/w) mixture of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀, Test Sample 2 was prepared in the same manner as the preparation of Sample 1 in Example 1.

10 Preparation of Sample 3

[0045] Using a 40:60 (w/w) mixture of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀, Test Sample 3 was prepared in the same manner as the preparation of Sample 1 in Example 1.

15 Preparation of Sample 4

[0046] Using a 60:40 (w/w) mixture of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀, Test Sample 4 was prepared in the same manner as the preparation of Sample 1 in Example 1.

20 Preparation of Comparative Sample 2

[0047] Using a 80:20 (w/w) mixture of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀, Comparative Sample 2 was prepared in the same manner as the preparation of Sample 1 in Example 1.

25 (2) Oral absorption test

[0048] Sample 1, Sample 2, Sample 3, Sample 4, Comparative Sample 1, and Comparative Sample 2 were used as test samples. The test was performed in the same manner as described in Example 1 except that the determination of total plasma coenzyme Q₁₀ concentration was carried out at 3 hours after administration.

30 [0049] The test results are presented in Fig. 2. In Fig. 2, the ordinate represents the total plasma coenzyme Q₁₀ concentration at 3 hours after administration and the abscissa represents the weight ratio of oxidized form of coenzyme Q₁₀ to reduced form of coenzyme Q₁₀ in the test sample. Each bar represents mean \pm standard deviation.

[0050] It is apparent from Fig. 2 that compared with the composition containing only the oxidized form of coenzyme Q₁₀ and the composition in which the reduced form of coenzyme Q₁₀ accounts for 20 weight % of total coenzyme Q₁₀, administration of the compositions in which reduced form of coenzyme Q₁₀ accounts for 40 weight % or more of total coenzyme Q₁₀ resulted in higher plasma coenzyme Q₁₀ concentrations. Moreover, with a weight ratio of contained reduced form of coenzyme Q₁₀ increasing, the plasma coenzyme Q₁₀ concentration was further increased. Those results indicate that because it contains 40 weight % or more of the reduced form of coenzyme Q₁₀, the medicinal composition of the present invention is absorbed in a definitely larger amount than the composition containing only the oxidized form of coenzyme Q₁₀ and the composition in which the reduced form of coenzyme Q₁₀ accounts for 20 weight % or less of the total coenzyme Q₁₀ content.

[0051] Then, using a 15:85 (w/w) mixture of oxidized form of coenzyme Q₁₀ and reduced form of coenzyme Q₁₀ (hereinafter referred to as main medicine) as an active ingredient, several dosage forms were prepared by the conventional pharmaceutical procedures.

45 Formulation Example 1 (powders)

[0052] The main medicine was dissolved in acetone and the solution was adsorbed on microcrystalline cellulose, followed by drying. The product was mixed with corn starch to provide powders in the routine manner.

Main medicine	10 Parts by weight
Microcrystalline cellulose	40 Parts by weight
Corn starch	55 Parts by weight

EP 0 956 854 A1

Formulation Example 2 (tablets)

[0053] The main medicine was dissolved in acetone and the solution was adsorbed on microcrystalline cellulose, followed by drying. The product was mixed with corn starch, lactose, carboxymethylcellulose, and magnesium stearate and the mixture was granulated in the routine manner by adding an aqueous solution of polyvinylpyrrolidone as a binder. To the granules thus obtained was added talc as a lubricant followed by mixing and the resulting composition was compressed into tablets each containing 20 mg of the main medicine.

Main medicine	20 Parts by weight
Corn starch	25 Parts by weight
Lactose	15 Parts by weight
Carboxymethylcellulose calcium	10 Parts by weight
Microcrystalline cellulose	40 Parts by weight
Polyvinylpyrrolidone	5 Parts by weight
Magnesium stearate	3 Parts by weight
Talc	10 Parts by weight

Formulation Example 3 (capsules)

[0054] The following components were granulated by the routine procedure and filled in hard gelatin capsule shells to provide capsules each containing 20 mg of the main medicine.

Main medicine	20 Parts by weight
Microcrystalline cellulose	40 Parts by weight
Corn starch	20 Parts by weight
Lactose	62 Parts by weight
Magnesium stearate	2 Parts by weight
Polyvinylpyrrolidone	3 Parts by weight

Formulation Example 4 (soft capsules)

[0055] Soybean oil was warmed to 60°C and the main medicine melted at 60°C was added and dissolved. Then, vitamin E was added gradually to prepare a homogeneous mixture, which was then processed into soft capsules each containing 20 mg of the main medicine.

Main medicine	20 Parts by weight
Vitamin E	15 Parts by weight
Soybean oil	350 Parts by weight

INDUSTRIAL AVAILABILITY


[0056] Because of the above constitution, the medicinal composition of the present invention is well absorbable after oral administration and shows a high level of bioavailability.

EP 0 956 854 A1**Claims**

1. A medicinal composition comprising coenzyme Q₁₀ as an active ingredient with the reduced form of coenzyme Q₁₀ accounting for more than 20 weight % of said coenzyme Q₁₀.
2. The medicinal composition according to Claim 1 wherein said reduced form of coenzyme Q₁₀ accounts for 40 weight % or more of the total coenzyme Q₁₀ content.


EP 0 956 854 A1

Fig. 1



EP 0 956 854 A1

Fig. 2



EP 0 956 854 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/02845

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl ⁶ A61K31/12, A61K31/05 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int. Cl ⁶ A61K31/12, A61K31/05 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS (STN), MEDLINE (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WEBER, C., et al., 'Effect of Deletary Coenzyme Q ₁₀ as an Antioxidant in Human Plasma.' Molec. Aspects. Med., (1994) Vol. 15 (Supplement), pages s97-s102	1 - 2
A	JP, 59-47202, A (Zeria Pharmaceutical Co., Ltd.), March 16, 1984 (16. 03. 84), Full descriptions (Family: none)	1 - 2
A	JP, 4-89456, A (Shionogi & Co., Ltd.), March 23, 1992 (23. 03. 92), Full descriptions (Family: none)	1 - 2
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document but published on or after the international filing date "C" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not to conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search November 12, 1997 (12. 11. 97)		Date of mailing of the international search report November 26, 1997 (26. 11. 97)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)